

# Keith Corpus, MD

## Shoulder Arthroscopy

---

<u>Postop</u>	<u>Goals</u>	<u>Precautions</u>	<u>Exercises</u>
<b>Weeks 0-4</b> PT 1-2x/week HEP daily	Edema and pain control FF 140°, ER 40° IR behind back	Sling 0-2 weeks comfort only DC sling over 2-4 weeks No abduction >90° No rotation in 90° abduction	Elbow, wrist, hand ROM Codman/Pendulum Shoulder P/AA/AROM as tolerated Isometrics (arm at side), begin at 2 weeks Posterior capsule stretch Posture training
<b>Weeks 4-8</b> PT 2-3x/week HEP daily	FF 160°, ER 60°	No Sling	Advance P/AA/AROM as tolerated Advance isometrics Progress to bands as tolerated
<b>Weeks 8-12</b> PT 2-3x/week HEP daily	Full ROM Return to activity	None	Continue ROM exercises Strengthening eccentric/closed chain
<b>Weeks 12+</b> HEP daily	No restrictions	None	Maintenance exercises

### AMENDMENTS TO PROTOCOL FOR CONCOMITANT PROCEDURES

- DISTAL CLAVICLE EXCISION:** Weeks 0-8: no cross-body adduction, abduction >90°, or rotation in 90°.
- BICEPS TENODESIS:** Weeks 0-4: sling while not doing exercises, no elbow flexion >1lb → Weeks 4-8: DC sling, begin biceps isometrics. → Weeks 8+: begin biceps resistance training.